

Detection and Classification of Alzheimer's Disease by Employing CNN

Smt. Swaroopa Shastri*

Department of Computer Science and Engineering (MCA), Visvesvaraya Technological University Kalaburagi, Karnataka

E-mail: swaroopas04@gmail.com

ORCID iD: <https://orcid.org/0000-0002-8897-5878>

*Corresponding author

Ambresh Bhadrashetty

Department of Computer Science and Engineering (MCA), Visvesvaraya Technological University Kalaburagi, Karnataka

E-mail: ambresh.bhadrashetty@gmail.com

ORCID iD: <https://orcid.org/0000-0003-4443-9018>

Supriya Kulkarni

Department of Computer Science and Engineering (MCA), Visvesvaraya Technological University Kalaburagi, Karnataka

E-mail: supriya191099@gmail.com

Received: 08 September 2022; Revised: 11 December 2022; Accepted: 13 January 2023; Published: 08 April 2023

Abstract: Alzheimer's illness is an ailment of mind which results in mental confusion, forgetfulness and many other mental problems. It effects physical health of a person too. When treating a patient with Alzheimer's disease, a proper diagnosis is crucial, especially into earlier phases of condition as when patients are informed of the risk of the disease, they can take preventative steps before irreparable brain damage occurs. The majority of machine detection techniques are constrained by congenital (present at birth) data, however numerous recent studies have used computers for Alzheimer's disease diagnosis. The first stages of Alzheimer's disease can be diagnosed, but illness itself cannot be predicted since prediction is only helpful before it really manifests. Alzheimer's has high risk symptoms that effects both physical and mental health of a patient. Risks include confusion, concentration difficulties and much more, so with such symptoms it becomes important to detect this disease at its early stages. Significance of detecting this disease is the patient gets a better chance of treatment and medication. Hence our research helps to detect the disease at its early stages. Particularly when used with brain MRI scans, deep learning has emerged as a popular tool for the early identification of AD. Here we are using a 12- layer CNN that has the layers four convolutional, two pooling, two flatten, one dense and three activation functions. As CNN is well-known for pattern detection and image processing, here, accuracy of our model is 97.80%.

Index Terms: Alzheimer's Disease (AD), Deep Structured Learning (DL), MRI.

1. Introduction

Computational neuroscientific approaches have been shown to be incredibly beneficial in far-reaching psychological wellbeing preliminary studies [1]. This interdisciplinary area of research may help simulate biological processes which regulate human mind in both its healthy & diseased phases, and it can also guide incorporation of these cycles into therapeutic practice. The rapid expansion of large-volume biological datasets (neuropsychology and allied organic information) during the last ten years, along with breakthroughs in AI (ML), has unlocked new paths for analysis and visualization of neurodegenerative & neuropsychiatric illnesses [2]. With computing standpoint, this new progress had resulted in evolution of instruments which include a few sufferer-specific perceptions into expectations and work on the therapeutic consequence of victim with such concerns [3,4]. People at high risk for Alzheimer's ill-health & AD-related mental effect are target population for such neuroscientific therapies, with hopes of improving underlying openness and completing treatment plan [5,6]. In light of above, ongoing research has focused on

developing extremely sophisticated approaches that employ machine learning frameworks to improve the gauging of Alzheimer's disease. Such treatments focus to develop underlying openness and treatment completion in individuals who are at danger for AD & AD-related mental decline [5,6] is the best existing solution. However, our paper provides a methodology that provides a greater variation in the output, it mentions different range in the disease after examining the MRI. Accuracy is one of the major problems in previous works, our paper intends to provide a solution that improves accuracy. This section provides an overview of the historical background of AD by summing together data about AD distributions obtained from a Google Scholar search. Only works that were disseminated relatively recently, between 2008 and 2019, were considered for inclusion. Previous experts' routines and strategies were taken into consideration. Kraepelin came up with the phrase "Alzheimer's sickness" while the clinical description of illness was still vague. Although the ultimate verdict on Auguste Deter sickness was not quite clear, a clinical diagnosis of Alzheimer's disease started to take shape that seemed realistic more than a century after its discovery. Alzheimer's disease, first illustrated Dr. Alois Alzheimer in 1907 & subsequently by Proskin in 1909, is characterized by presence of amyloid plaques and neurofibrillary tangles. Researchers from Munich University, Germany and MaxPlanck Institute of Neurobiology, Martinsried discovered in 1998 that neurofibrillary spasms and amyloid plaques can affect specific cerebrum parts. Since then, this investigation has been regarded as the first publicized example of Alzheimer's disease; more importantly, the case fits current models of how the disease is defined. Figure1 is a side-by-side contrast of healthy brain & affected by Alzheimer's disease. As of right now, promotion ranks as the sixth leading reason of mortality in United States. Moreover, current evaluations suggest that issue may attempt to take 3rd place as top cause of mortality among aged. Alzheimer's disease is a devastating illness, so it's obvious that early detection and treatment are top priorities. Diagnosing Alzheimer's disease requires battery of medical exams & mountain of detailed information. In cases when brain problem can be predicted with reasonable accuracy, ordering an MRI test may be useful treatment option, but it can also be rather intimidating. All the research mentioned up until now in the paper have been done using many methodologies but lack in accuracy which has been covered by our paper. Research objectives include detecting the disease in early stages using CNN and classifying it into no, mild, moderate and severe dementia. The restrictions of our offered model are that it performs a little poorly on negative images. The flow of the paper is as follows, segment 2 gives information on literature survey, segment 3 is about proposed methodology used, dataset, results and discussion and segment 4 contains conclusion.




Fig.1. Progress of AD from Mild Cognitive Impairment to utmost AD.




Fig.2. Structure of a 12-layer CNN.

Fig.1. shows a pictorial representation of a healthy and an alzheimer's effected brain and the progress of brain from healthy to severe AD effected brain. Fig.2. represents the layers of a 12- layer CNN. CNNs, are one subset of CBNs. In 2019, the Alzheimer's Association described that there is still no medical care for Alzheimer's sickness. Over 50,00,000 Americans are incapacitated by Alzheimer's illness, with two hundred thousand of those affected are younger than 65. By 2050, 10 million Americans, mostly adults over the age of 60, are projected to have Alzheimer's disease, according to the research. Mind structure may be gleaned from underlying imagery., together with neurons, synapses, glial cells, and other structures. Utilitarian imaging provides information on the cerebrum's activities. The structural MRI, that assesses mind vol into vivo to modify cerebrum declension, is the most frequently used MRI for AD occurrences (deprivation of tissue, cells, neurons, and so on.). Degeneration of the mind is an unavoidable moderate

aspect of Alzheimer's disease. A sample of a mock MRI used to show brain decay is shown in Fig.3. In Fig.4, we see schematic of fMRI, common way for analysing the prefrontal cortex and other parts of the brain. For example, Figure 5a depicts regions of brains of aged individuals (AD sufferer and controls), and Figure 5b depicts the typical transitory behavior of a group with comparable criteria.




Fig.3. Instance of chief MRI.





Fig.4. Instance of utilitarian Magnetic Resonance Imaging (fMRI).



(a)



(b)

Fig.5. (a) Cerebrum area into more seasoned controls & AD; (b) MRI filter mind in cathartic Temporal decline.

Studies have shown that SPECT can accurately predict cerebral perfusion when employed in the examination of patients suspected of having Alzheimer's disease. A recent work examined 116 Alzheimer's patients. There were 23 age-matched controls, 26 individuals having non-dementia Alzheimer's, & 67 participants with other neurological diseases. The participants were split into groups with dementia and those without it. 99mTCHMPAO CSF-tau peptide volume were linked to SPECT processing. The factors chosen improved the precision of the assessment, resulting in a trustworthy evaluation. Other studies focused on parietal and transitory trauma in Alzheimer's victims, and they discovered substantial links between their neuropsychological test results and SPECT ends. SPECT is more useful for assessing AD contrasted with CSF-tau protein. A primitive blood vessel turns name perfusion picture of a 82-year-old patient is shown in Fig5. The contiguous segment of the canal tree is not recommended for this kind of evaluation because of the severe stall arterial transitime.

- PET: One imaging technique that uses radioactive atoms to map workings of brain is known as positron emission tomography (PET). Amyloid and fluorodeoxyglucose are two most often used tracers in the diagnosis of AD, as seen in

Figure 7. Observing, focusing one's attention, contemplating, recalling, and IGUE are all included.




Fig.6. Basic proposition of autostrada spin labelling.




Fig.7. PET of brain in typical circumstances.

Radioligands C-PMP & C-MP4A have been employed to identify acetylcholinesterase. And if this data is to be believed, AD sufferers' temporal lobes have contracted. Similar decline was seen in subjects with MCI who eventually acquired AD. Patients with Alzheimer's disease and neurodegenerative dementia were further divided into groups. Because those with AD showed gains compared to those with Parkinson's disease and frontotemporal lobar degeneration (FTLD) (PD), Pittsburgh substance B 11C-PIB) were used as A-beta amyloid-specific ligands. In contrast, since they allow researchers to see differences in nigrostriatal dopaminergic neurons, neuroreceptors & FP-CITSPECT are much valuable & practical. For water diffusion analysis, FP-CIT SPECT imaging is used. This technique can be used to determine the location, directional, & anisotropy of white matter into brain. DTI is not regarded as a reliable technique for evaluating CSF biomarkers despite substantial research into the CSF-tau biomarker & amyloid stages. Fig.8. shows the DTI for Alzheimer's illness done with tractography examination, which shows methodical alteration.




Fig.8. DTI in Alzheimer's illness.




Fig.9. Illustration of Convolutional Neural Network (CNN).

To attain the optimum goal, used and minimized descriptors employ DL. As a result, as shown in Fig.9, the CNN is utilized to get the depictions from the pictures. CNNs are very good in retrieving nonexclusive attributes. Regardless, the provided images do not reflect the progressions in the photo dataset. As a result, different strategies are used depending on the problem. Pre-existing DL models, such as CNNs, may be enhanced with the help of a fresh dataset in order to take on a different kind of problem.

2. Related Works

[1] Bullmore and Sporns worked on complex brain structures and analysis of brain on how a brain functions and its network through their work "Analyzing the anatomical and functional systems of the brain using graph theory". [2] Eickhoff with the help of others, MRI neuroimaging, both functional and structural, is now the gold standard for studying human brain structure in vivo. To evaluate task-evoked variations on neurogenesis as in framework of cognitive neuroscience investigations, magnetic resonance imaging (MRI) has been widely used since the 1990s. [6] L. Zhang al. al"attention deficit hyperactivity disorder 3D CNN Based Automatic Diagnosis., 2003, "Understanding Morphological and Anatomical Fmri." In drawing on the research of physicians, L. Zhang his coworkers created a three-dimensional Network approach to analyze the regional spatial patterns of MRI data using the standard way to analyze headpictures.[7] keuck worked on rediscoveries of old stuff like historical patient files and samples of brain images of firstcasesofAD.[8] In contrast to various cancers, AD patients would eventually lose their cognitive function, talents, etc K. Trojachanec et al. Using a range of time series, longitudinal brain MRI retrieval for Hypertension K. Trojachanec and associates discussed the initiatives taken to enhance Hypertension case report recovery. [9] In their article "Towards Alzheimer's disease classification by transfer learning," authors M. Hon , N. M. Khan provide new insight into the use of neuroanatomical data in the detection of Alzheimer's disorder.[10] Adlard and Sundberg described the increasing advancementinchemistry.[11] Swaroopa Shastri and A Patange once a restorative man needs to concentrate a patient's therapeutic history; once proposal and counter letters are composed for a patient minded by complex facilities; while patient is in crisis and in this way the medicinal comfort history wishes to be surveyed. [12] Aptly titled "An ANN model for diagnostic scoring predictions in Alzheimer disease utilizing structure neuroimaging measurements," N bhagwat's effort is really laudable. [13] J. Yao, "Improvement of a multimodal system for cardiovascular figured tomographygating," sheds light on few important topics regarding alzheimer's. [3] Chaddad and others worked on "Profound radiomic investigation of x-ray connected with alzheimer's infection,". [4] Weiming Guo and colleagues proposed an approach that uses the Alzheimer's Disease Neuroimaging Project's standardised MRI datasets. This technique achieves an accuracy of 79.9% in verifying exit crossings, with an AUC of 86.1%. [14] The study of N. Hill, J. Mogle, "Alzheimer's disease risk variables as mediator of subjective memory loss & objective memory decline: methodology for construct-level replica analysis," focuses upon that memory difficulties an AD victim might encounter and the construct-level replica analysis.[15] The study of K. F. Tracker and coworkers, Management of Fecal Incontinence in Frail Older Adults Community-based evidence on the impact of fecal incontinence on daily life. [5] S. Ahmed and others have shared their work regarding "Troupes of fix based classifiers for determination of alzheimer illnesses," that explains importance of classification and the classifiers used in AD detection Recent Evaluations of the Earliest Cases of Hypertension," by L. Keuck et al. [16] The main subject of this book is the accounts of medical scientists who have discovered and revised outdated information. The 1990s saw a Hamburg psychotherapist and a Bavarian molecular neuropsychiatrist in command of the artifacts search. R. H. Blank et al. Choosing Final Decisions in Individuals with Adhd Across Cultures In this essay, a number of moral dilemmas surrounding judgments about edge care for Dementia sufferers are covered. It concentrates on subjects like physician-assisted suicide, advance directives, diet, antibiotics, and other forms of hospital attention. They went into the Mri data.[17] They also applied D. Jin et allongitudinal .'s Attention-based 3D CNN for particular level and diagnosing Hypertension , which was reported in Nature [18] Jee-young Han and others work on Alzheimer's suggested that cholinesterase inhibitor may not improve cognitive course in MCI. [19] Vatanabe, Manzine and others described the tale of Alzheimer's. Their work also included biomarkers from recent area of diagnosis. [20] Emtiaz Hasan and colleagues at Bg Universities in Nigeria investigated potential analysis of Alzheimer's disease utilising 12-layer CNN and compared it to an 8 layer CNN utilising Functional mri also as basis. Weiming Guo et colleagues. reported Brain mri processing using dcnns for the prediction of Alzheimer's disease from moderate memory loss. [21] Raigonda, M. R., Terdal, S. P., & Raigond, B. (2022). Detection of the viral disease on the potato foliar and tubers using a machine learning approach. International Journal of Health Sciences. By adopting the new technology and continuous monitoring, the diseases can be identified at the initial stage.[22] Rani, Megha & Terdal, Sujatha. Design Engineering A Review on the Disease Identification on the Potato Foliar and Tuber. Texture defines the orientation of the form. Variations in brightness or grey level values are the roughness or bumpiness in this instance. These are the many texture attributes that need to be evaluated for categorization purposes.

3. Methods

A. Proposed Methodology




Fig.10. Proposed model.




Fig.11. Methodology used.

In this section, we are discussing our proposed model that is the 12-layer CNN model that involves the following steps as shown in Fig.11. The block diagram shows seven basic steps in our proposed method.

Step1: Dataset collection – We have collected our data from OASIS dataset that stands for Open Access Series of Imaging Studies. The dataset contains MRI of both men and women.

Step2: Data pre-processing - The dataset contains photos of various sizes. The architecture may be less accurate due to the various image sizes. We pre-process data for this reason. Image denoising is a fundamental problem in image processing and computer vision. To improve the performance of our model, we denoise.

Step3: Image segmentation- Often based on the properties of the picture's pixels, image segmentation is a widely used method in digital image processing and analysis to divide an image into various parts or areas. Foreground and background can be distinguished in an image by segmenting it, or pixels can be grouped together based on their similarity in colour or shape. Here MRI is divided into various parts so that analysis can be easier.

Step4: Feature extraction- By choosing and combining variables into features, feature extraction helps to extract the best feature from those large data sets, effectively lowering the amount of data. These features are simple to use

while still accurately and uniquely describing the real data set.

Step5: Data split- Usually, data splitting is done to prevent overfitting. In that situation, a machine learning model fits its training data too well and is unable to consistently fit new data.

Step6: Model evaluation- Model evaluation is the practise of employing several evaluation measures to comprehend the performance and strengths and weaknesses of a machine learning model. With the early stages of research, it is crucial to evaluate a model's effectiveness. Model evaluation also aids in model monitoring. Evaluating our proposed model by giving different MRI images to know its efficiency.

Step7: Classification- Finally we classify the results that go through the model as no, mildly, moderate, severe dementia.

The suggested model's accuracy, precision is compared to those of many existing state-of-the-art CNN models using OASIS dataset. The principal result of this work is a 12- layer convolutional neural network (CNN) models whose accuracy of 97.80% exceeds that of any previously available CNN model on this dataset. Here, we discuss 12-layer CNN model we've built for Alzheimer's disease categorization using brain MRIs. Convolutional layer selection, layer pooling selection, layer flattening, & full activation of associated layers and functions are the five stages that make up our CNN 12-layer model. Our model has four sublayers of convolution, a pooling layer between each layer of convolution, flattening the layer using the two-layer dense layer that we are utilising here, and three activation functions to demonstrate our model's performance. CNN provides filters that make distinctions based on edge and colour and bend. MRI image is fed as the input to the model it goes through the above mentioned layers. Convolutional layer performs convolutional operation on the input fed to it and is passed to next layer. Next step is the pooling layer helps to lessen the output from the previous layer which is called downsampling by sliding a filter on it. Flatten layer flattens the network i.e it makes any multidimensional input into one dimensional. Activation function is used here to make easier for the model to understand complex patterns in the input provided.

B. Dataset

Thanks to the OASIS datasets hosted on central.xnat.org, a user-friendly platform for studying cognitive, clinical, & neuroimaging facets of normal aging is available to community at no cost. These datasets span a broad variety of demographic, cognitive, & genetic features. All of our information comes from the OASIS database. Open Access Series of Imaging Studies abbreviates to OASIS. A total of 410 people were included in this sample for the cross-sectional analysis. The ages of these participants range from 17 to 95 years. On view in Figure 9 are a few examples of brain MRIs that may be found as in Open Access Series of Imaging Studies dataset. To put it another way, eighty percent of the data is used for training, while the persist twenty percent is employed for testing.




Fig.12. Sample pictures from Open Access Series of Imaging Studies database.

CNN model on this dataset. Here, we discuss 12-layer CNN model we've built for Alzheimer's disease categorization using brain MRIs. Convolutional layer selection, layer pooling selection, layer flattening, & full activation of associated layers and functions are the five stages that make up our CNN 12-layer model. Our model has four sublayers of convolution, a pooling layer between each layer of convolution, flattening the layer using the two-layer dense layer that we are utilising here, and three activation functions to demonstrate our model's performance. CNN provides filters that make distinctions based on edge and colour and bend.

C. Results and Discussion




Fig.13. Accuracy graph of 12- layer CNN.

The above given graph clearly shows how the proposed model is better than the previous model used, with proposed model having accuracy of 97.80%. The proposed model has four main layers and several sublayers of CNN which makes it total of twelve layers providing higher accuracy and precision.

4. Conclusions

Alzheimer's is infirmity that effects both physical and mental health that is why it is important to detect it as early as possible which is done satisfactorily by our model. A high-level literature evaluation led us to the conclusion that our proposed system that uses twelve-layer CNN has outdone the previous systems with higher accuracy. The work is well recognised and well welcomed, but because the majority of the patients investigated have the condition, it doesn't contribute much to the early diagnosis of Alzheimer's disease. This study examined technique and recognition using a sampling of the most significant AD datasets. We have used this framework to compare with different CNN architectures. This technique can be used for brain imaging research at an early stage.

References

- [1] E. Bullmore and O. Sporns, "Complex mind organizations: Graph hypothetical examination of primary and utilitarian frameworks," *Nature surveys. Neuroscience*, vol. 10, pp. 186-98, 03 2009.
- [2] S. Eickhoff, T. E. Nichols, J. D. V. Horn, and J. A. Turner, "Sharing the abundance: Neuroimaging information stores," *NeuroImage*, vol. 124, pp. 1065 - 1068, 2016.
- [3] A. Chaddad, C. Desrosiers, and T. Niazi, "Profound radiomic investigation of x-ray connected with alzheimer's infection," *IEEE Access*, vol. 6, pp. 58213-58221, 2018.
- [4] Weiming Lin, Tong Tong, Qinquan Gao, Di Guo, Xiaofeng Du, Yonggui Yang, Gang Guo, Min Xiao, Min Du, Xiaobo Qu, Alzheimer's Disease Neuroimaging Initiative, "Convolutional Neural Networks-Based MRI Image Analysis for the Alzheimer's Disease Prediction from Mild Cognitive Impairment", 2018 Nov 5;12:777. doi: 10.3389/fnins.2018.00777. eCollection 2018.
- [5] S. Ahmed, K. Y. Choi, J. J. Lee, B. C. Kim, G.- R. Kwon, K. H. Lee, and H. Y. Jung, "Troupes of fix based classifiers for determination of alzheimer illnesses," *IEEE Access*, vol. 7, pp. 73373-73383, 2019.
- [6] L. Zou, J. Zheng, C. Miao, M. J. Mckeown and Z. J. Wang, "3D CNN Based Automatic Diagnosis of Attention Deficit Hyperactivity Disorder Using Functional and Structural MRI," in *IEEE Access*, vol. 5, pp. 23626- 23636, 2017.
- [7] L. Keuck, "History as a biomedical matter: recent assessments of the first cases of alzheimer's disease," *History and Philosophy of the Life Sciences*, vol. 40, no. 1, p. 10, 2017.
- [8] K. Trojchanec, I. Kitanovski, I. Dimitrovski, and S. Loshkovska, "Longitudinal brain mri retrieval for alzheimer's disease using different temporal information," *IEEE Access*, vol. 6, pp. 9703-9712, 2017.
- [9] M. Hon and N. M. Khan, "Towards Alzheimer's disease classification through transfer learning," in *Proc. IEEE International Conference on Bioinformatics and Biomedicine (BIBM)*, Kansas City, pp. 1166-1169, 2017.
- [10] E. R. Adlard, "Richard j. sundberg: The chemical century. molecular manipulation and its impact on the 20th century," *Chromatographia*, vol. 80, no. 10, pp. 1599-1599, 2017.
- [11] Swaroopa Shastri, A Patange "International Journal of Engineering Science" vol.7, Issue 7, Pp.14272-14275, 2017
- [12] N. Bhagwat, "Prognostic applications for alzheimer's sickness utilizing attractive reverberation imaging and AI," Ph.D. exposition, Graduate Department of Institute of Biomaterials and Biomedical Engineering University of Toronto, 2018.
- [13] J. Yao, "Improvement of a multimodal system for cardiovascular figured tomography gating," Ph.D. dissertation, School of Electrical and Computer Engineering, Georgia Institute of Technology, 2018.
- [14] N. Hill and J. Mogle, "Alzheimer's disease risk factors as mediators of subjective memory impairment and objective memory decline: protocol for a construct-level replication analysis," *BMC Geriatrics*, vol. 18, 2018.
- [15] K. F. Tracker, M. Northwood, V. Haggard, and F. Bates, *Management of Fecal Incontinence in Frail Older Adults Living in the Community*. Springer International Publishing, 2018, pp. 127-148.
- [16] R. H. Blank, *End-of-Life Decision Making for Alzheimer's Disease Across Cultures*. Springer Singapore, 2019, pp. 121-136.
- [17] D. Jin, J. Xu, K. Zhao, F. Hu, Z. Yang, B. Liu, T. Jiang, and Y. Liu, "Attention-based 3d convolutional network for alzheimer's disease diagnosis and biomarkers exploration," in *2019 IEEE 16th International Symposium on Biomedical Imaging (ISBI2019)*. IEEE, 2019, pp.1047- 1051.
- [18] J.- Y. Han, L. Besser, C. Xiong, W. Kukull, and J. Morris, "Cholinesterase inhibitors may not benefit gentle mental hindrance and gentle alzheimer diseasedementia," *Alzheimer disease and associated disorders*, vol.33, pp. 87-94, 04 2019.
- [19] I. Vatanabe, P. Manzine, and M. Cominetti, "Historic concepts of dementia and alzheimer's disease: From ancient times to the present," *Revue Neurologique*, 2019.
- [20] Emtiaz Hussain, Mahmudul Hasan, Syed Zafrul Hassan, Tanzina Hassan Azmi, Md Anisur Rahman, Mohammad Zavid Parvez, "Deep Learning Based Binary Classification of Alzhiemer's Disease Detection using Brain MRI Images", 978-1-7281-5169- 4/20/\$31.00 c 2020 IEEE.
- [21] Raigonda, M. R., Terdal, S. P., & Raigond, B. (2022). Detection of the viral disease on the potato foliar and tubers using a machine learning approach. *International Journal of Health Sciences*, 6(S4), 9336-9354. <https://doi.org/10.53730/ijhs.v6nS4.10740>.
- [22] Rani, Megha & Terdal, Sujatha. (2022). Design Engineering A Review on the Disease Identification on the Potato Foliar and Tuber.

Authors' Profiles



Ms. Swaroopa Shastri working as an Assistant Professor in the Department of Computer Science and Engineering (MCA) at Visvesvaraya Technological University, Centre for PG Studies, Kalaburagi. She is in the field of Computer Applications at VTU's, CPGS, Kalaburagi, Karnataka, India. She is in teaching profession for more than 10 years. She has presented/published 20 papers in National and International Journals and Conference. Her main area of interest includes Machine Learning, Image Processing and Artificial Intelligence.



Mr. Ambresh Bhadrashetty working as an Assistant Professor in the Department of Computer Science and Engineering(MCA) at Visvesvaraya Technological University, Centre for PG Studies, Kalaburagi. He is in the field of Computer Applications at VTU's, CPGS, Kalaburagi, Karnataka, India. He is in teaching profession for more than 10 years. He has presented/published 20 papers in National and International Journals and Conference. His main area of interest includes Biomedical Data Mining, Machine Learning and Artificial Intelligence.

How to cite this paper: Smt. Swaroopa Shastri, Ambresh Bhadrashetty, Supriya Kulkarni, "Detection and Classification of Alzheimer's Disease by Employing CNN", International Journal of Intelligent Systems and Applications(IJISA), Vol.15, No.2, pp.14-22, 2023. DOI:10.5815/ijisa.2023.02.02